

Factorization and Lie Point Symmetries of General Lienard-type Equation in the Complex Plane

Özlem Yeşiltas^{†‡1}

[†]*Department of Mathematics and Statistics, Concordia University 1455 de Maisonneuve Boulevard West, Montreal Quebec, Canada H3G 1M8*

[‡]*Department of Physics, Faculty of Arts and Sciences, Gazi University, 06500 Ankara, Turkey*

Abstract

We present a variational approach to general Lienard type equation in order to linearize it and as an example Van der Pol oscillator is discussed. The new equation which is almost linear is factorized. The point symmetries of the deformed equation is also discussed and the two dimensional Lie algebraic generators are obtained.

Key words: Lienard, symmetry

¹e-mail : yesiltas@gazi.edu.tr

1 Introduction

Investigation of partial and ordinary nonlinear differential equations is one of the popular recent problems in mathematical physics. A typical example of a relaxation oscillator is known as Van der Pol limit cycle [1]. Van der Pol equation can be adapted to study the Lienard equation. This paper mainly devoted to study Lienard type equations whose dynamic properties have been widely studied due to the applications in many fields such as physics, mechanics and engineering [1, 2, 3, 4, 5, 6, 7, 8]. This kind of general equations and their properties are studied within many methods. Factorization method [9] is one of these topics and it can be applied to a class of nonlinear differential equations including commonly Lienard type equations [10, 11, 12]. Beyond the factorization method, Lie group approach which is an integrability method is a powerful tool for obtaining solutions and investigating the symmetry properties of the differential equations [13, 14, 15]. In this paper, variational approach and factorization of the system are discussed in section 2. In addition, section 3 involves a short discussion of the Lie symmetries of the system, the results are concluded in section 4.

2 Linearizing and factorization

During the process of deriving Euler's equation one writes

$$y(x, \alpha) = y(x, 0) + \alpha y_1(x). \quad (1)$$

If we write a difference equation of the form

$$\Delta f = f[x, y(x, \alpha), y'(x, \alpha)] - f[x, y(x, 0), y'(x, 0)] \quad (2)$$

$$= \left(\frac{\partial f}{\partial y} y_1 + \frac{\partial f}{\partial y'} y_1' \right) \alpha + \dots \quad (3)$$

and then continue with introducing the variation of f as

$$\delta f = \frac{\partial f}{\partial y} \delta y + \frac{\partial f}{\partial y'} \delta y' \quad (4)$$

where $\delta y = \alpha y_1$ and $\delta y' = \alpha y_1'$. Thus the variation of f is expressed in terms of constant α . Thus, this operation can be applied to a differential equation called as general Lienard equation:

$$\ddot{x} + F(x(t))\dot{x} + x = 0 \quad (5)$$

where the dots correspond derivatives of $x(t)$ with respect to t . Now apply the variational operator to (5) that is defined by $\delta x^n = nx^{n-1}\delta x = y^n$. Here the differential and variational operators are commuting such that $\delta \frac{dx}{dt} = \frac{d}{dt} \delta x$. In this condition (5) becomes

$$\ddot{y} + F(x(t))\dot{y} + \delta(F(x(t)))\dot{x} + G(t)y = 0. \quad (6)$$

As an example, we can discuss Van der Pol type equation where $F(x(t)) = \varepsilon(1 - x(t)^2)$. In this case (5) will be introduced as a complex differential equation [16]

$$\ddot{z} - \varepsilon(1 - |z|^2)\dot{z} + z = 0 \quad (7)$$

where the complex variable $z = z_1 + iz_2$ and the equation admits the translation symmetry $z(t) \rightarrow z(t + \delta)$, phase symmetry $z(t) \rightarrow e^{i\theta} z(t)$, spatial symmetry $z(t) \rightarrow -z(t)$, conjugation symmetry $z(t) \rightarrow z^*(t)$. (7) can be written in a general form:

$$\ddot{z} - \varepsilon(1 - |z|^2)\dot{z} + G(t)z = 0. \quad (8)$$

where $G(t)$ is an unknown function. If the variational operation is applied to (8)

$$\delta \ddot{z} - \varepsilon(1 - |z|^2)\delta \dot{z} + \varepsilon \dot{z} \delta |z|^2 + G(t) \delta z = 0 \quad (9)$$

is obtained and (8) turns into the form

$$\ddot{Y} + \varepsilon(|z(t)|^2 - 1)\dot{Y} + \varepsilon\dot{z}(t)Y^2 + G(t)Y = 0 \quad (10)$$

here $\delta z_1 = Y(t)$ is used, we assume that $\delta z_2 = 0$ in the equation given above, according to non-analytical zz^* . Henceforth, we can discuss the factorization of (10). We use a homographic transformation $D(\alpha, \beta, \theta)$ that is generally given by

$$(y, t) \rightarrow (Y, T) : y = \frac{\alpha(t)Y + \tilde{\beta}(t)}{\gamma(t)Y + \delta(t)}, \quad \alpha\delta - \tilde{\beta}\gamma \neq 0, \quad T = \theta(t), \quad \tilde{\beta}(t) = \beta_0\beta(t) \quad (11)$$

and makes the ordinary differential equation (10) is form invariant. Under this transformation $D(\alpha, \beta, \theta)$, (10) turns into

$$\begin{aligned} & \frac{d^2Y}{dT^2} + \frac{1}{\theta'} \left(2\frac{\alpha'}{\alpha} + \frac{\theta''}{\theta} - \varepsilon(x^2 - 1) \right) \frac{dY}{dT} \\ & - \varepsilon\dot{x} \frac{\alpha}{\theta'^2} Y^2 + \left(\frac{\alpha''}{\alpha} + 2\varepsilon\dot{x}\tilde{\beta} - \varepsilon(1 - x^2) \frac{\alpha'}{\alpha} \right) \frac{Y}{\theta'^2} \\ & + \frac{1}{\alpha^{-1}\theta'^2} (\tilde{\beta}'' + \varepsilon\dot{x}\tilde{\beta}^2 - \varepsilon(1 - x^2)\tilde{\beta}' - G(t)) = 0, \quad \alpha\theta' \neq 0 \end{aligned} \quad (12)$$

In order to transform (12) into an equation below that is addressed as canonic form of type I in [17]:

$$\frac{d^2Y}{dT^2} - 6Y^2 + 6\beta_0 = 0 \quad (13)$$

one must write

$$\begin{aligned} \theta'^2 &= -\frac{\varepsilon\dot{x}\alpha}{6} \\ (Ln \ \alpha)' &= -\frac{2}{5}(\varepsilon(1 - x^2) + \frac{\ddot{x}}{2\dot{x}}) \\ \tilde{\beta} &= -\frac{1}{2\varepsilon\dot{x}}(-G(t) + \varepsilon(1 - x^2)(Ln \ \alpha)' + (Ln \ \alpha)'' + (Ln \ \alpha')^2). \end{aligned} \quad (14)$$

In [18] it is pointed out that (14) is essential to obtain (13) and equations in (14) must be satisfied. In this study we will follow a more general way, there won't be such a restriction given in (14). We will use

$F_1(t) = \varepsilon(|z(t)|^2 - 1)$, $F_2(t) = \varepsilon\dot{z}(t)$ in (10) that can be factorized as

$$\left(\frac{d}{dt} - f_2(Y)\right)\left(\frac{d}{dt} - f_1(Y)\right)Y = 0 \quad (15)$$

where f_1 and f_2 satisfy the relations:

$$F_1 = -\left(f_1 + f_2 + Y\frac{df_1}{dY}\right) \quad (16)$$

$$F_2Y + G = f_1f_2.$$

We want to show that if (10) admits the factorization for f_1 and f_2 have the form

$$f_1 = \alpha(\sqrt{F_2Y} + i\sqrt{G}) \quad (17)$$

$$f_2 = \alpha^{-1}(\sqrt{F_2Y} - i\sqrt{G})$$

where α is a constant and can be found as $\alpha = \pm i\sqrt{\frac{2}{3}}$ by comparing (17) and (16). Then one can find a relation between F_1 and G using the first equation in (16)

$$F_1 = \pm \frac{5}{\sqrt{6}}\sqrt{G}. \quad (18)$$

The second bracket in (15) allows us to write

$$\dot{Y} \mp i\sqrt{\frac{2F_2}{3}}Y^{\frac{3}{2}} \pm \sqrt{\frac{2G}{3}}Y = 0 \quad (19)$$

which is a Bernoulli differential equation and we obtain the solutions of the form [19]

$$Y = -\mu(t)^2 \left(\int^t \sqrt{\frac{F_2(t')}{6}} \mu(t') dt' \right)^{-2}, \quad \mu(t) = e^{\mp \int^t \sqrt{\frac{G(t')}{6}} dt'} \quad (20)$$

In order to derive special exact solutions, assume that $(\partial_t - f_1)Y = \psi(t)$, and it yields

$$\dot{Y} - f_1(Y)Y = \psi(t) \quad (21)$$

$$\dot{\psi} - f_2(Y)\psi = 0 \quad (22)$$

and this choice will lead to have more general solutions. In [10] the author selected a constant function $f_2(Y)$ but we want to use a non-constant f_2 which is given by the second formula in (17). Then, (21) can be given by

$$\dot{Y} = f_1(Y)Y + e^{\int^t f_2(Y)dt'}. \quad (23)$$

Thus, Y can be derived easily as solving this equation

$$\dot{\omega} \mp \sqrt{\frac{G(t)}{6}}\omega = \mp i\sqrt{\frac{F_2}{6}} - \frac{1}{2}\omega^3 e^{\mp i\sqrt{\frac{3}{2}}\int^t \left(\sqrt{\frac{F_2}{\omega}} - i\sqrt{G}\right)dt'}, \quad \omega = Y^{-\frac{1}{2}}. \quad (24)$$

The solution is introduced as [19]

$$\omega(t) = \pm \frac{i}{\mu(t)} \int^t \sqrt{\frac{F_2(t')}{6}} \mu(t') dt' - \frac{1}{2\mu(t)} \int^t \mu(t') \varphi(t') e^{\mp \int^t \sqrt{\frac{3}{2}}G(t')dt'} \quad (25)$$

where t dependent function φ is defined by

$$\frac{\dot{\varphi}}{\varphi} = \mp i\sqrt{\frac{3F_2(t)}{2\omega(t)}} + \frac{3\dot{\omega}}{\omega}. \quad (26)$$

Assume that φ is a constant. Thus, a special solution for ω is given by

$$\omega(t) = \frac{1}{4} \left(C_1 \pm i \int^t \sqrt{\frac{F_2(t')}{6}} dt' \right)^2 \quad (27)$$

where C_1 is an integration constant. Now, let us choose F_2 as

$$F_2(t) = -6e^{\mp \sqrt{6}\int^t \sqrt{G}dt'} \quad (28)$$

then, using (28) and (18), the following expression can be obtained

$$\dot{z} = -\frac{6}{\varepsilon} e^{\mp \frac{6\varepsilon}{6}\int^t (|z|^2 - 1)dt'} \quad (29)$$

and the solution of z is introduced by

$$z = ce^{i\theta t} \quad (30)$$

where c and θ are some constants. In this case, (29) is satisfied when

$$\varepsilon = i \quad (31)$$

$$c(c^2 - 1) = \mp 5 \quad (32)$$

$$\theta = \pm \frac{1 - c^2}{5}. \quad (33)$$

Thus, F_1 , F_2 and G are expressed by

$$F_1 = i(c^2 - 1) \quad (34)$$

$$F_2 = -c\theta e^{i\theta t}$$

$$G = \frac{6}{25}(1 - c^2).$$

3 Symmetries of the deformed equation

The point symmetry of (10) will be discussed here. As it is stated in [11], the differential equations that forms a system

$$F_i(x, u, u_{(1)}, \dots, u_{(k)}) = 0, \quad i = 1, \dots, s \quad (35)$$

is invariant under the X which is an infinitesimal generator such that

$$XF_i|_{F_i=0} = 0. \quad (36)$$

So, the infinitesimal generator for (10) will have a form

$$X = \xi(x, y) \frac{\partial}{\partial x} + \eta(x, y) \frac{\partial}{\partial y} \quad (37)$$

for a second order equation

$$y'' = f(x, y, y') \quad (38)$$

where the point transformations in the plane are given by

$$\bar{x} \approx x + a\xi(x, y) \quad (39)$$

$$\bar{y} \approx y + a\eta(x, y) \quad (40)$$

Thus we have

$$X(y'' - f(x, y, y'))|_{y''=f} = 0. \quad (41)$$

The equation above can also be written [13]

$$\begin{aligned} & \eta_{xx} + (2\eta_{xy} - \xi_{xx})y' + (\eta_{yy} - 2\xi_{xy})y'^2 - y'^3\xi_{yy} - \xi f_x - \eta f_y \\ & + (\eta_y - 2\xi_x - 3y'\xi_y)f - (\eta_x + (\eta_y - \xi_x)y' - y'^2\xi_y)f_{y'} = 0 \end{aligned} \quad (42)$$

In this case, $f = -F_1y' - F_2y^2 - Gy$ and here the derivatives are taken with respect to x and y is used instead of Y here. We can group the equations as

$$(y')^3 : \xi_{yy} = 0 \quad (43)$$

$$(y')^2 : \eta_{yy} - 2\xi_{xy} + 2F_1\xi_y = 0 \quad (44)$$

$$y' : 2\eta_{xy} - \xi_{xx} + \xi F_{1x} + F_1\xi_x = 0 \quad (45)$$

$$y^2 : \xi F_{2x} - F_2(\eta_y - 2\xi_x) = 0 \quad (46)$$

$$y : \xi G_x + 2F_2\eta - G(\eta_y - 2\xi_x) = 0 \quad (47)$$

$$y^0 : \eta_{xx} + \eta_x F_1 - \eta G = 0. \quad (48)$$

$$yy' : 3G\xi_y = 0 \quad (49)$$

Thus we can obtain ξ from (43)

$$\xi = \alpha(x)y + \beta(x). \quad (50)$$

On the other hand, from (44), η takes the form

$$\eta = W(x)y + S(x). \quad (51)$$

$W(x) = 0$ and $\alpha(x) = 0$ can be obtained by using (47) and (49). If $\eta_y - 2\xi_x$ in (46) is used in (47) we get

$$2F_2\eta + \beta(x)\frac{F_{2x}}{F_2}G = 0 \quad (52)$$

and using (45)

$$\beta(x) = A + Be^{F_1x} \quad (53)$$

where A, B are constants. We can obtain $S(x)$ using (48) as

$$S(x) = C_1e^{-\alpha_+x} + C_2e^{-\alpha_-x} \quad (54)$$

where C_1, C_2 are constants and $\alpha_{\pm}$ can be given in terms of the parameters of (10):

$$\alpha_{\pm} = \frac{1}{2} \left(F_1 \pm \frac{\sqrt{F_1^2 + 4G}}{2} \right). \quad (55)$$

If we use the values of F_1, F_2 and G and (47) we have

$$2F_2\eta - G(\eta_y - 2\xi_x) = 0 \quad (56)$$

where the derivative of G is zero as it is seen from (34). Then, we can obtain

$$\alpha_+ = -i\theta + F_1, \quad \alpha_- = i\theta \quad (57)$$

Finally we get the value of θ which is same with the result obtained in the previous section

$$\theta = \frac{1 - c^2}{5} \quad (58)$$




Figure 1: The skeleton of (10) for some specific parameters. $\dot{Y} = P$, $\ddot{Y} = Q$. $\mathcal{M} = \{Y, \dot{Y}, \ddot{Y}\}$ reduced manifold.

which means that the positive sign should be used for θ in (33). Thus, two dimensional generators are obtained as

$$X_1 = \frac{\partial}{\partial x} + e^{-\alpha_+ x} \frac{\partial}{\partial y} \quad (59)$$

$$X_2 = e^{F_1 x} \frac{\partial}{\partial x} + e^{-\alpha_- x} \frac{\partial}{\partial y} \quad (60)$$

the commutation between these operators is $[X_1, X_2] \neq 0$ and the group is not closed.

4 Conclusion

The flexibility of the variational approach is shown by applying the variational operator to the Lienard type equation and linearized Van der Pol differential equation is examined in the complex plane. The deformed differential equation is factorized and solutions are obtained. Moreover, it is shown that the

time dependent coefficients of the linearized equation that are, F_1 and F_2 , found as pure imaginary and complex for the real values of c . Thus, interesting result is the parameter ε is obtained as i . It means that the symmetry properties of the equation can be changed under the time reversal symmetry within the contributions of F_1 , F_2 and G such that the original equation (7) is invariant under spatial reversal but (10) is not invariant under $Y \rightarrow -Y$ and it is not invariant under both spatial and $\varepsilon = i$, $i \rightarrow -i$ parameter and $t \rightarrow -t$ reversal. Besides factorization, the symmetry of equation (10) is examined within point symmetries and two dimensional generators are obtained. The group is not closed, however we have concluded the parameter θ of $z(t)$ is the same as that obtained with both the symmetry equations and factorization results.

The skeleton for (10) is a flat surface such that this equation has two-dimensional generators, as is shown in figure 1.

References

- [1] T A Burton, Stability and Periodic Solutions of Ordinary and Functional Differential Equations, Orland: Academic Press, 1985; A H Nayfeh, Introduction to Perturbation Techniques, John Wiley and Sons Inc, 1981; A A Andronov, A A Vitt, S E Khaikin , Theory of oscillators, Dover, New York, 1989.
- [2] J K Hale, Theory of Functional Differential Equations, New York: Springer-Verlag, 1977.
- [3] Zhou Jin, Sun Shu, Liu Zengrong, Appl. Math Comput, 2005, 161(2) 655-666.
- [4] H A Antosiewicz, J London Math Soc, 1955, 30 64-67.
- [5] H I Freedman, Y Kuang, Uniqueness of limit cycles in Lienard-type equation, Nonlinear Anal, 1990, 15 333-338.
- [6] G Sansone, R Conti, Non-Linear Differential Equation, New York: MacMillan, 1964.
- [7] M M Kamel, Physica Scripta, 79(2) 2009 025004.
- [8] Zhang Hua-Yan, Ran Zheng, Chinese Physics Letters, 26(3) 2009 030203.
- [9] L Infeld, T E Hull, Rev. Mod. Phys. 23 1951 21.
- [10] O Cornejo-Perez, J. of Phys. A: Math. and Gen., 42(3) 2009 035204.
- [11] O Cornejo-Perez, H C Rosu, Progress of Theo. Phys., 114(3) 2005 533-38.
- [12] H C Rosu, O Cornejo-Perez, Phys. Rev. E 71 2005 046607.

- [13] N H Ibragimov, Elementary Lie Group Analysis and Ordinary Differential Equations, J Wiley and Sons Ltd., 1999.
- [14] R K Gazizov and C M Khalique, Hindawi Publishing Corporation Mathematical Problems in Engineering Volume 2006, Article ID 68753, Pages 111 DOI 10.1155/MPE/2006/68753.
- [15] A Karasu Kalkanlı, P G L Leach, Journal of Math. Phys. 50(7) (2009).
- [16] R López-Ruiz and Y Pomeau, Phys. Rev. E 55 1997 R3820.
- [17] E L Ince, Ordinary Differential Equations, Ordinary Differential Equations, 1956 (New York-Dover).
- [18] P G Estvez, Ş Kuru, J Negro and L M Nieto, J. Phys. A: Math. Theor. 40 2007 9819.
- [19] E Kreyszig, Advanced Engineering Mathematics, Wiley, ISBN 0471488852 (2005).